
## **CORY'S SHEARWATER**

Calonectris diomedea borealis



## Regional Occurrence (BCRs 14, 30)

- Distribution: pelagic in region, generally July-October
- Habitats used: pelagic; breeding in burrows and crevices on islands

NAWCP ND=2

## **Regional Abundance**

- 1980s population size: 10,000-100,000nb
- Historical population size: unknown
- % change in regional population: unknown

NAWCP PT=3

## **Regional Threats**

- Identified threats: Habitat loss by development of coastal area
- Introduced predators in breeding areas

NAWCP TN=3

Conservation Status: Abundance (b=breeding, nb=nonbreeding, t=total individuals) and Conservation Rank

Global	North America	BCR 14	BCR 30	Mid-Atlantic	S. New England	Gulf of Maine	Maritimes
?	?	?nb	?nb	?nb	?nb	?nb	?nb
IUCN-	NAWCP-	Moderate	Moderate	NJ- SZN; DE-SZN	NY- S?N; CT- no info	MA-SN; NH- SZN; ME- no	QC, PE- no info
Not Evaluated	Moderate Concern	(PI=?)	(PI=?)	MD-SZN; VA-SN	RI- SZN; MA-SN; VT-no info	info; NB, NS- no info	NB, NS- no info

**Conservation Needs**: Sustainability:

Density and Foraging Habitat. 30-100 birds/ sq km; feeds at or near the surface as secondary or tertiary predators on fish, larvae, crustaceans; frequently follows fishing boats; *Issues*: ; *Population Goals*: ; *Habitat Goals*: